

Chateau Drug

451 4th Street, Ketchum, Idaho 83340 ph 208-726-2596
fax 208-726-8268

Patient Name: _____ Address: _____

DOB: _____ Allergies: _____ Phone number: _____

Compound drug : _____ Amount: _____ Refills: _____

Directions: _____

Doctor name: _____ Address: _____

Phone number: _____ DEA# _____ NPI: _____

Doctor signature: _____ Date: _____

Female Hormones

- Biestrogen capsule: Ratio: 80/20 70/30 50/50 Dose: 0.625mg
1.25 mg 2.5 mg or _____ mg
- Progesterone: SR capsule 25 mg 50mg 100 mg 200 mg or _____ mg
- Progesterone troche: 50 mg 100 mg or _____ mg
- Testosterone capsule: 1 mg 2 mg 4 mg or _____ mg
- Estriol 0.5% vaginal cream
- Progesterone Cream 5% 10% 20% (circle one)

Amount: _____ Refills: _____

Directions: _____

Non-hormonal libido enhancer

- 2% Sildenafil 3% Aminophylline 6% Arginine Cream

Amount: _____ Refills: _____ Apply ¼ gram of cream to clitoris
and external vaginal area 30 minutes prior to sexual activity.

Male Hormones

- Testosterone 200 mg/ml cream
- Testosterone 50 mg/ml solution
- Testosterone Gel: 1% 2.0 % 2.5% (circle one)

Amount: _____ Refills: _____

Directions: _____

Topical Pain Creams & Gels

- 2% Diclofenac 3% Baclofen 2% Cyclobenzaprine 2% Gabapentin 6% Tetracaine cream

Amount: _____ Refills: _____

Directions: _____

Wound and Scar Topical Creams & Gels

- 0.1% Fluticasone, 8% Levocetirizine 3% Pentoxifyline 15% Ubiquinol in Anhydrous Cream

Amount: _____ Refills: _____

Directions: _____

Psychological Topical Transdermal Creams Gels

- Quetiapine, venlafaxine, trazodone, buspirone, escitalopram, lorazepam (circle one)

Drug: _____ % in PLO transdermal gel Amount: _____ Refills: _____

Directions: Apply 1 or 2 grams to inner arm; rub into skin. Wash hands.

Male ED Oral Troches (aka Lozenges or capsules)

- Sildenafil 27 mg 55 mg 110 mg
- Tadalafil 6 mg 12 mg 22 mg
- Vardenafil 6 mg 12 mg 22 mg

Amount: _____ Refills: _____

Directions: Lozenges: Dissolve one troche between cheek and gum 30 minutes prior to activity. Capsules: Take 1 capsule orally ½ hour prior to activity for erectile dysfunction.

Hospice Formulation 2% Lidocaine 1% Hydrocortisone suppository

Amount: _____ Refills: _____ Directions: Insert 1 suppository rectally 2-4 times daily

Antiviral Formulation

- 5% Acyclovir 10% Benzocaine 2% Sunscreen in lip balm Amount: _____
Refills: _____
- 5% Acyclovir/5% Lidocaine Lip Balm Amount: _____ Refills: _____

Directions: Apply topically to lips 4-5 times daily after cold sore outbreak for 5-10 days

- 0.2 % Deoxy-d-glucose solution (nasal spray) Amount: _____ Refills: _____

Directions: Apply 1-2 sprays in each nares three times daily.

Atypical

- Low dose Naltrexone 4 mg capsule Amount: _____ Refills: _____

Directions: Take 1 capsule at bedtime daily for pain relief.